

***DURIO TANJUNGPURENSIS* (MALVACEAE), A NEW SPECIES AND ITS ONE NEW VARIETY FROM WEST KALIMANTAN, INDONESIA**

ZIDNI ILMAN NAVIA AND TATIK CHIKMAWATI*

*Department of Biology, Bogor Agricultural University, Darmaga Campus,
Bogor, 16680, Indonesia*

Key words: *Durio tanjungpurensis*, *D. tanjungpurensis* var. *tembaganensis*, *D. testudinarum*, new species, new variety, West Kalimantan

Abstract

A new species and a new variety of *Durio* Adanson from West Kalimantan are described namely, *Durio tanjungpurensis* Z.I. Navia, **sp. nov.** and *D. tanjungpurensis* var. *tembaganensis* Navia, **var. nov.** The new species has small fruits (5 - 8 cm in diam), brown to yellowish green pericarp, milky white thin aril (<1 mm), and not edible. It is related to *D. testudinarum* Becc. *Durio tanjungpurensis* varies in its flower and fruit characters. The new variety differed from *D. tanjungpurensis* Z.I. Navia, **sp. nov.** in six characters, i.e. leaf ovate, petal oblanceolate with apex round, fruits on the branches, on the stem, at the base of trunk, no-spine area in fruit apex, pericarp green to yellowish green, aril only 20 - 40% covering the seeds.

Introduction

Durio Adanson (1958) is the largest genus in the Family *Malvaceae* (Nyffeler and Baum 2001) comprising of 30 species which are confined to the tropics of Southeast Asia. Its centre of distribution is Borneo Island, Indonesia, and 21 species of which 15 are endemic. Further, 11 species are distributed in Malaya, 8 species in Sumatra, and one species in each in Myanmar (Burma) and Srilanka (Kostermans 1958, Uji 2005, Mansur 2007). The genus *Durio* is characterized by five diagnostic features, i.e. a densely lepidote or stellate indumentum on the lower side of the leaves on young branches, and various part of the flowers; a cup-shaped calyx; rather large, showy flowers with generally whitish or reddish petals; anthers consisting of clusters of globular or elongate pollen locules which open either via an apical pore or a longitudinal slit; and big spiny fruits with large seeds that are covered with a fleshy or, in some species, rather leathery aril (Nyffeler and Baum 2001). Eight *Durio* species have edible fruits including *D. zibethinus* Murr., *D. kutejensis* (Hassk.) Becc., *D. graveolens* Becc., *D. dulcis* Becc., *D. oxleyanus* Griff., *D. testudinarum* Becc., *D. lowianus* Scort. ex King, and *D. grandiflorus* (Mast.) Kosterm. et Soeng. (Soengeng 1962, Salma 2011).

Materials and Methods

In a recent taxonomic study of *Durio*, survey for specimen collection was conducted in 10 districts of West Kalimantan, Indonesia, namely Mempawah, Landak, Bengkayang, Sanggau, Sekadau, Sintang, Kapuas Hulu, Melawi, Kayong Utara and Ketapang. The survey uncovered occurrence of a new species in two districts i.e., Sekadau and Kapuas Hulu. This species is very similar to *D. testudinarum* Becc.

After a number of literature study (Bakhuizen van den Brink 1924, Kostermans and Soengeng 1958, Kostermans 1958, Salma 2011) and examinations carried out on a number of specimens, it is concluded that these plants actually represent one new species namely, *Durio tanjungpurensis* Z.I. Navia, **sp. nov.** and a new variety namely *Durio tanjungpurensis* var. *tembaganensis* Navia, **var. nov.**

*Author for correspondence: <tchikmawati@yahoo.com>.

Results and Discussion

As the newly reported species *Durio tanjungpurensis* has some similarity with *D. testudinarum*, a comparison of characters has been presented in Table 1. From Table 1 it is observed that *D. tanjungpurensis* differs distinctly from *D. testudinarum* Becc over six diagnostic characters and the species hitherto described as a new species of the Genus *Durio*. A taxonomic detail of the newly described species has been furnished below.

Table 1. Morphological comparison of *D. testudinarum* and *D. tanjungpurensis*.

Characters	<i>D. testudinarum</i>	<i>D. tanjungpurensis</i>
Aril	2 - 3 mm, consumed	<1 mm
Colour of aril	White to yellow	Milky white
Favour of aril	Sweet and watery	Plain
Colour of fruit	Yellowish-brown to brownish-green	Yellowish-green to dark brown
Bud apex	Pointed and curved	Round
Status	Cultivated	Wild

***Durio tanjungpurensis* Z.I. Navia, sp. nov.**

(Figs 1, 2)

Diagnosis: *Durio tanjungpurensis* est morphologia similis ad *D. testudinarum*, sed ita habet parvus fructus (5 - 8 cm in diam), brunneus pericarpium, lactaneus arillus tenues (<1 mm), et ita est fructus est non edibilis.

Durio tanjungpurensis is morphologically similar to *D. testudinarum*, but it has small fruit (5 - 8 cm in diam), brown pericarp, milky white thin aril (<1 mm), and its fruit is not edible.

Holotypus: West Kalimantan Province: Sekadau Regency, Rawak Subdistrict, Rawak forest, 0°09'15.8"S - 110°51'14.8"E, 45 meter a.s.l., April 2013, Z.I. Navia 25.

Tree, 10 - 20 m tall, 14 - 32 cm in diam. Outer bark brown to dark brown; inner bark red brown. Sapwood white. Leaves simple, alternate; petiole 10- 23 mm long; leaf blade thinly to thickly leathery, oblong to elliptic, obovate lanceolate, 12 - 44 × 3 - 12.5 cm; base acute to round; apex acuminate, long acuminate, caudate, acumen about 5-20 mm long; upper surface glabrous, midrib and primary nerves sunken; lower surface densely covered with tightly appressed, midrib strongly prominent, lateral nerves parallel, reticulation inconspicuous. Inflorescences solitary or 2 - 5 flowers clustered in a group on the trunk, at the base of trunk; flower buds green to greenish yellow, ellipsoid, 2 - 3 × 1 - 2 cm, apex acute to round; pedicel 3 - 5.5 cm long; epicalix persistent, splitting into 2 equal, ovate, 2 - 3.5 × 1.5 - 2.5 cm, outside densely scaly, inside stellate-hairy; calyx 5, free, green, ovate, apex acute, 2 - 3 × 1 - 1.5 cm, outside scaly, inside glabrous with dense stellate hairs at the base; petal 5, free, white, oblong, apex round to acute, 5.6 - 7.5 × 1 - 1.5 cm, outside dense stellate hairs, thickened at middle of base; stamens whitish yellow, shorter than the corolla, upper 5 - 10 mm of phalanges; filaments free, slender, bearing each with 10 reniform anthers in a globose clump; anthers opening by a slit; ovary densely covered with scales; stigma yellow, capitate, 5-lobed, glabrous. Fruit on the trunk, at the base of trunk, oblate to globose, lobed; 6.7 - 7.8 cm in diam.; spines conical and pyramidal, 5 - 11 mm long; aril very thin <1 mm, aril milky white.




Fig. 1. *Durio tanjungpurensis* **sp. nov.** A. Tree and fruits, B. Leaves, C. Bud, D. Flowers and buds, E. Bud, F. Anther and stigma, G. Ovary covered with scales, H. Aril milky white, I. Fruits at the base of trunk and above ground level.

Phenology: Flowering in May - September, Fruiting in November - December.

Distribution and habitat: *Durio tanjungpurensis* had been found in five locations in West Kalimantan, that were Rawak forest, Rejunak forest, Merindang Hill, Tembaga forest and Sagu Hill. It grows in mixed Dipterocarp forest, customary forest communities, near plantation; associated to *Artocarpus*, *D. griffithii*, *D. kutejensis*, *D. oxleyanus*, *D. zibethinus*, *Hevea braziliensis*, *Lithocarpus*, *Shorea*, *Ficus*, *Nepenthes*, *Rattan*, *Pandanaceae*.




Fig. 2. Illustration of *Durio tanjungpurensis* sp. nov. A. Leaves, B. Flower, C. Petal, D. Anther, E. Pistil, G. Ovary, H. fruit, I. Fruit apex with spines.

Taxonomic relationships: *Durio tanjungpurensis* is morphologically similar to *D. testudinarum* Becc. It has leaf oblong; leaf apex acuminate to long acuminate, base acute, rounded to cordate, and has fruit on the base of trunk.

Etymology: Named after the type locality, West Kalimantan

Additional specimen examined: INDONESIA. West Kalimantan Province, Sekadau Regency, Nanga Mahap, Tembaga forest, *Z.I. Navia* 01, *Z.I. Navia* 02, *Z.I. Navia* 03, *Z.I. Navia* 04, *Z.I. Navia* 05, *Z.I. Navia* 06, *Z.I. Navia* 07, *Z.I. Navia* 08; Nanga Engkulun Subdistrict, Merindang Hill, *Z.I. Navia* 09, *Z.I. Navia* 10, *Z.I. Navia* 11, *Z.I. Navia* 12, *Z.I. Navia* 13, *Z.I. Navia* 14; Rawak Subdistrict, Rejunak forest, *Z.I. Navia* 16, *Z.I. Navia* 17, *Z.I. Navia* 18, *Z.I. Navia* 19, *Z.I. Navia* 20, *Z.I. Navia* 21, *Z.I. Navia* 22, *Z.I. Navia* 23; Rawak forest, *Z.I. Navia* 25, *Z.I. Navia* 26, *Z.I. Navia* 27; Kapuas Hulu Regency, Nanga Silat Subdistrict, Sagu Hill, *Z.I. Navia* 28, *Z.I. Navia* 29, *Z.I. Navia* 30, *Z.I. Navia* 31, *Z.I. Navia* 32, *Z.I. Navia* 33, *Z.I. Navia* 34, *Z.I. Navia* 35.

Note: There is variation in flower and aril, shape, size and color of fruit among the specimens of *D. tanjungpurensis*. The first group of specimens grows flowers and fruits at the base of trunk, fruits with brown pericarp and aril completely covering the seeds. The author named the first group as var. *tanjungpurensis*. The second group has flowers and fruits on the branches, at the base of stem and at the base of trunk, green pericarp, aril very thin only 20-40% covering the seeds. The author named the second group as var. *tembaganensis* var. nov.

Key to varieties

Petal oblong, fruits brown pericarp, conical spines, aril completely covering the seed

var. **tanjungpurensis**

Petal oblanceolate, fruits green to yellowish green pericarp, pyramidal spines, aril 20-40% covering the seeds

var. **tembaganensis** var. nov

Durio tanjungpurensis var. **tanjungpurensis**

(Figs 3, 4)

Holotype: West Kalimantan Province: Sekadau Regency, Rawak Subdistrict, Rawak forest, 0°09'15.8"S - 110°51'14.8"E, 45 meter a.s.l., April 2013, *Z.I. Navia* 25.

Petiole 10 - 23 mm long; Leaf thinly to thickly leathery; oblong to elliptic, obovate-lanceolate, 12 - 44 × 3 - 12,5 cm; petal oblong, apex acute to round; fruit on the trunk, at the base of trunk, oblate to globose, lobed; brown pericarp until ripe; 6.7 - 7.8 cm diam.; spines conical, 5 - 11 mm long; aril very thin <1 mm, milky white aril, covered seed completely.

Distribution: it has been found in four locations in West Kalimantan, that are Rawak forest, Rejunak forest, Merindang Hill, and Sagu Hill.

Durio tanjungpurensis var. **tembaganensis** Navia, var. nov.

(Fig. 2)

Diagnosis: *Foliis ovatus, petalum oblanceolatus cum apex rotundus; fructus in ramiflorus, in caulibus, ad basim e truncus, spineless area in fructus apex grandis, pericarpium viride to flavo-vireus, arillus 20-40% tegens at semen.*

Holotype: West Kalimantan Province: Sekadau Regency, Nanga Mahap Subdistrict, Tembaga forest, 0°30'37.4"S - 110°40'48.9"E, 93 meter a.s.l., April 2013, *Z.I. Navia* 04.

Leaves ovate; petal oblanceolate with apex round; fruits on the branches, on the stem, at the base of trunk, no-spine area in fruit apex large, pericarp green to yellowish green, aril 20 - 40% covering the seeds.



Fig. 3. *Durio tanjungpurensis* var. *tembaganensis* var. nov. A. Tree, B. Flowers, C. Bud with apex rounded, D. Pistil, E-G. Fruits on the branches, on the stem and at the base of trunk, H. Fruit apex without spines. I-J. Fruit with aril not completely covering the seed.

Petiole 11 - 13 mm long; Leves thinly; oblong to elliptic, ovate, 12 - 44 × 3 - 12,5 cm; petal oblanceolate, apex round, 6.6 - 7.7 × 1.2 - 1.5 cm; fruit on the branches, on the stem, at the base of trunk, globose, lobed; pericarp green to yellowish green; 6.3 - 7.8 cm diam.; spines pyramidal, 3 - 5 mm long, no-spine area in fruit apex large, aril very thin, aril milky white, aril 20 - 40% covering the seeds.

Distribution and habitat: *Durio tanjungpurensis* var. *tembaganensis* var. nov has been found only in one location, Tembaga forest in West Kalimantan. It grows in mixed Dipterocarp forest, customary forest communities, near plantation society; associated to *Artocarpus*, *D. kutejensis*, *D. oxleyanus*, *D. zibethinus*, *Hevea braziliensis*, *Lithocarpus*, *Shorea*, *Ficus*, *Rattan*, *Pandanaceae*.




Fig. 4. Illustration of *Durio tanjungpurensis* var. *tembaganensis* var. nov. A. Leaves, B. Flower, C. Petal, D. Anther, E. Pistil, F. Sepal, G. Ovary, H. Fruit, I. Fruit apex without spines.

Acknowledgements

The authors are thankful to Prof. Mien A. Rifai for his guidance and advice during this research. Thanks are due also to Adi Bejo Suwardi, M.Si for his assistance during exploration in the study area.

References

- Bakhuizen van den Brink RC 1924. Revisio Bombacaceae. Bulletin J. Bot. Buitenzorg, Series III **6**(2): 162-163, 223-225.
- Kostermans AJGH 1958. The genus *Durio* Adans. (*Bombacaceae*). Reinwardtia **4**(3): 47-153.
- Kostermans AJGH, Soegeng WR 1958. A monograph of the Genus *Durio*. Commun. For. Res. Inst. Bogor **61**: 1-51
- Mansur M 2007 Penelitian ekologi jenis-jenis durian (*Durio* spp.) di desa Intuh Lingau Kalimantan Timur. J. Tek Ling **8**(3): 211-216.
- Nyffeler R, Baum DA 2001. Systematics and character evolution in *Durio* s. lat. (*Malvaceae/Helicteroideae/Durioneae* or *Bombacaceae-Durioneae*). Organisms Divers. Evol. **1**: 165-178.
- Salma I 2011. *Durio of Malaysia*. Mardi, Malaysia. pp. 117-125.
- Soegeng WR 1962, The species of *Durio* with edible fruits, Economy Bot. **16**: 270-282
- Uji T 2005. Keanekaragaman jenis dan sumber plasma nutfah *Durio* (*Durio* spp.) di Indonesia. Buletin Plasma Nutfah. **11**: 28-33.

(Manuscript received on 3 March, 2015; revised on 11 May, 2015)